

Effects of Climate Change on Local Marine, Estuarine, and Riverine Fishes



Several climate-related changes have the potential to affect critical fish habitat as well as abundance and distribution of fish on the Oregon coast.

- *Changes to marine habitat may reduce biodiversity and alter the distribution of fishes.*
- *Degradation and loss of estuarine habitat may jeopardize the reproductive success of local fish.*
- *Alterations to stream hydrology may result in critical habitat loss for cold water species.*



Salmon spawning habitat
Photo: Umpqua Watersheds

Pacific storm
Photo: Partnership for Coastal Watersheds

Climate-related changes such as sea level rise (SLR), ocean acidification, and increasing ocean temperature are expected to affect marine and estuarine fish habitat. Other changes such as altered precipitation patterns and increased frequency and severity of flood and drought, are expected to affect freshwater fish habitat. Fish that inhabit a variety of environments at different life stages, such as anadromous salmonids, are likely to be affected by all climate-related changes that affect both marine and estuarine habitat as well as freshwater habitat.

Sea Level Rise (SLR)

Many fish species use estuarine and near-shore ocean habitats at various parts of their life stages. Pacific herring (*Clupea pallasii*), for example, require eelgrass beds, rocky shorelines or other substrates on which to attach their eggs during breeding season (Monaco and Emmett 1990). Many species of forage fish vital to the marine food web, such as surf smelt (*Hypomesus pretiosus*) and sand lance (*Ammodytes hexapterus*), use estuaries as breeding areas (Glick et al. 2007). Estuaries are vital to anadromous species by providing rearing habitat, the availability and quality of which affects

Background

Sea Level Rise: *Our local NOAA tide station in Charleston has documented an average rate of sea level rise (SLR) of 1.3 mm (0.05 inches) over the past 30 years (0.42 feet in 100 years)(NOAA Tides and Currents 2006). The rate of SLR is expected to accelerate over time. For example, according to the National Research Council (2012), predicted SLR rates for the area to the north of California's Cape Mendocino (the study's closest site to the Coos estuary), are reported as high as +23 cm (9 inches) by 2030; +48 cm (19 inches) by 2050; and +143 cm (56 inches) by 2100 .*

ocean survival in salmon (Miller and Simenstad 1997).

The availability of high quality estuarine habitat may be threatened by SLR. Scientists have not yet determined whether sand flat and mudflat elevations relative to tidal levels will be able to keep pace with SLR. In other words, sedimentation rates may adjust with sea level rise so sand and mud flats remain about the same elevation relative to tidal levels as they are now, resulting in very little change in fish habitat availability. However, it is unlikely that intertidal habitats will be allowed to migrate inland due to the high value of real estate near the existing high tide zone (Glick et al. 2007; Yamanaka et al. 2013). As SLR threatens rocky, intertidal habitat, the availability of hard substrate for egg deposition may decline.

Species diversity and distribution may be affected by SLR in instances where salt water encroaches on brackish and freshwater habitats. Glick et al. 2007 note that since aquatic animals have specific salinity tolerances, SLR-driven salinity changes will be beneficial for some species and unfavorable for others. They also suggest SLR may still affect fishes less sensitive to salinity changes because their food sources may be affected by changing salinity regimes even if they are not.

Ocean Acidification (OA)

Few studies have been conducted investigating the effect of OA on fish in temperate marine ecosystems (Ishimatsu et al. 2004). However, a growing body of research suggests that OA causes a wide range of deleterious physiological responses in marine fishes. Ishimatsu et al. (2004) explain that elevated levels of ambient CO₂ is associated with a condition in fish known as "hypercapnia," which causes disturbances that limit the function of the respiratory, circulatory, and nervous systems in fish. They suggest that the long-term effects of hypercapnia may inhibit important life functions by reducing growth, reproduction, and calcification.

Scientists who are studying tropical ecosystems report that OA may have significant effects on tropical fish. Dixon et al. (2012), Devine (2012) and Munday (2009) found significant effects of OA on the development of sensory mechanisms in tropical fish, and report that exposure to acidified seawater may impair the ability of these fish to recognize olfactory clues necessary for predator avoidance in tropical reefs.

Studies have shown that the effects of exposure to elevated CO₂ levels is greatest fish eggs, larvae, and juveniles, suggesting that fish in early developmental stages may be the most vulnerable to the impacts of ocean acidification (Kikkawa et al. 2003; Ishimatsu et al. 2004).

Other studies suggest that OA may change important fish habitat. Palacios and Zimmerman (2007) found that higher CO₂ concentrations are positively correlated with reproductive output, below-ground biomass, and vegetative proliferation in eelgrass (*Zostera marina*). However, they note that this response is not necessarily beneficial to fish that are associated with eelgrass meadows, because other characteristics of CO₂-rich environments (e.g., prolific algae growth and diminished water quality) are likely to overwhelm the positive effects of increased eelgrass productivity.

Although the precise effect of acidification on local fish populations is uncertain, it's likely that ocean acidification would reduce marine biodiversity through the loss of pH and CO₂-sensitive species and the likely reduction of habitat complexity (Widdicombe and Spicer 2008).

Increasing Ocean Temperature

Increasing ocean temperatures may affect the distribution of marine and estuarine fish, with warmer temperatures favoring habitats closer to the poles and nearer to the bottom of the ocean (Perry et al. 2005). Radovich (1961) has documented this phenomenon on the Pacific coast of North America by correlating unusually warm sea-surface temperatures with an increased number of anomalous fish landings between 1957 and

Background

Ocean Acidification: *Since the late 18th century, the average open ocean surface pH levels worldwide have decreased by about 0.1 pH units, a decrease of pH from about 8.2 before the industrial revolution to about 8.1 today (Feely et al. 2010). A 0.1 change in pH is significant since it represents about a 30 percent increase in ocean acidity (the pH scale is logarithmic, meaning that for every one point change in pH, the actual concentration changes by a factor of ten). Scientists estimate that by 2100 ocean waters could be nearly 150% more acidic than they are now, resulting in ocean acidity not experienced on earth in 20 million years (NOAA PMEL Carbon Program 2013). The best Pacific Northwest ocean acidification data we have so far is from the Puget Sound area where pH has decreased about as much as the worldwide average (a decrease ranging from 0.05 to 0.15 units)(Feely et al. 2010).*

1959. He cites several instances of warm-water species being caught north of their expected ranges including the following: Pacific bonito (*Sarda chiliensis*) in Eureka, California, skipjack tuna (*Katsuwonus pelamis*) off Cape Blanco in Oregon, swordfish (*Xiphias gladius*) in Monterey Bay, and dolphinfishes (*Coryphaena spp.*) as far north as Grays Harbor, Washington.

Background

Increasing Ocean Temperatures:

Worldwide, ocean temperatures rose at an average rate of 0.13°F per decade between 1901 to 2012. Since 1880, when reliable ocean temperature observations first began, there have been no periods with higher ocean temperatures than those during the period from 1982 – 2012. The periods between 1910 and 1940 (after a cooling period between 1880 and 1910), and 1970 and the present are the periods within which ocean temperatures have mainly increased (EPA 2013b).

Describing how the worldwide trend translates to ocean temperature trends off the Oregon coast is a complicated matter because of the high variability of sea surface temperatures influenced by coastal upwelling processes influenced by multiple climatic events that occur in irregular cycles (e.g., El Nino events). We do have 27 years (1967-1994) of water temperature data collected from near the mouth of the Coos estuary that indicate through preliminary analyses a very weak trend towards warming water temperatures (SSNERR 2013). Fifteen years (1995-2010) of data from multiple stations further up the South Slough estuary show very little water temperature change (Cornu et al. 2012).

Perry et al. (2005) note that fish with slower developmental rates or more complex life histories are less capable of adjusting to warming temperatures through rapid demographic responses like movement towards the poles. They anticipate that fish with these characteristics are more likely to be affected by rising ocean temperatures, because of their inability to rapidly respond to unfavorable habitat changes.

In addition to distributional responses, increased temperatures may affect fish by encumbering basic life functions. The amount of energy allocated toward growth and reproduction in fish usually declines as temperatures approach the extreme ends of species-specific tolerance ranges (Roessig 2004). This is demonstrated in the English sole, which has demonstrated significantly slower growth in temperatures above 17.5 °C (63.5 °F) and is likely to experience reduced growth in extreme estuarine temperatures (Yoklavich 1982; Rooper 2003). Similarly, studies of sand lances (*Ammodytes* spp) indicate that water temperature plays an important role in spawning timing and affects both recruitment and growth rates (Monaco and Emmett 1990).

Increased water temperatures also have the effect of compromising habitat quality for cold water species. According to ODFW (2014), higher water temperatures may accelerate the loss of areas that provide important cool water refugia and resting habitat for anadromous salmonid species.

Marine and Estuarine Hypoxia

Most of the climate-related alterations to fish habitat suggest increased likelihood of

Background

Local Effects of Changing

Ocean Conditions: *The physical conditions of an estuary are sensitive to changes in long-term oceanographic fluctuations. O'Higgins and Rumrill (2007) studied the physical response of the South Slough to changes in the PDO index by monitoring water quality in the South Slough estuary from 2000 to 2006. Their data show a positive and statistically significant relationship between temperature and the PDO index. They also found a negative and statistically significant relationship between dissolved oxygen and the PDO index. This suggests that local estuaries are both anomalously warm and less oxygenated during the warmer (positive) phases of the PDO. Similarly, Hamilton (2011) has studied the relationship between the physical conditions of local waters and ENSO events between 2004 and 2010. Her data demonstrate a positive and statistically significant relationship between temperature and a multivariate ENSO index at stations in Charleston, South Slough's Valino Island, and South Slough's Winchester Creek.*

secondary effects. For example, the potential loss of tidal marshes can lead to reduced water quality in estuaries, because tidal marshes regulate nutrients and filter

pollutants (Glick et al. 2007). High nutrient levels combined with increasing ocean temperatures and adequate light provide the ideal conditions for explosive algae growth. The over production of algae can damage aquatic ecosystems by blocking sunlight and reducing oxygen levels in the water column (EPA 2013a). Hypoxia, low levels of dissolved oxygen in water, has deleterious effects on fish. Hypoxic conditions are linked to limited reproductive function in several species of marine and estuarine fish (Giorgi and Congleton 1984; Landry et al. 2007; Thomas et al. 2007). In extreme circumstances, low oxygen levels have caused mass mortalities (Pacific Fishery Management Council 1983).

Changing Ocean Conditions

Climate change is likely to cause changes in a variety of ocean conditions which will affect fish:

- Climate change will likely affect ocean circulation and have some effect on Pacific coast upwelling patterns (Hayward 1997; Bakun 1990).
- Research suggests a correlation between ocean temperature and increased severity and frequency of storms (Knutson et al. 2010; Webster et al. 2005; McGabe et al. 2001).
- The oceanographic effects of climate change may directly affect the abundance and distribution of marine fishes by affecting the availability of food resources. For example, Monaco and Emmett (1990) found that food availability for larval northern anchovy (*Engraulis mordax*) is reduced by

storms or strong upwelling conditions. However, they also find that storms increase food abundance for adults.

- ODFW (2014) suggests that rising temperature may be cause for increased ocean stratification, a trend which has previously been associated with poor foraging conditions for salmonids.

In addition to the effects of climate change, Pacific Decadal Oscillation (PDO) and El Niño Southern Oscillation (ENSO) are cyclical patterns of climate variability (see sidebar next page) that influence ocean conditions,

Background

Pacific Decadal Oscillation and El

Niño Southern Oscillation: *The Pacific*

Decadal Oscillation (PDO) and El Niño Southern Oscillation (ENSO) are cyclical climatic patterns that affect weather and ocean currents in and around the Pacific ocean. PDO is a pattern of oceanic conditions that shifts every few decades. During a cold (negative) phase, the west Pacific warms, and the east Pacific cools; the opposite is true of a warm (positive) phase. ENSO is a climatic event that tends to occur every two to seven years and is characterized by anomalous warming of tropical Pacific waters (Mysak 1986).

Locally, this warming is associated with drier conditions, warmer temperatures, and lower precipitation and streamflow, although it can also result in greater winter storminess and flooding.

and fish abundance and distribution. The ecological response to shifts in PDO first affects primary producers and consumers before working to higher level consumers such as salmon. Warm PDO periods may be associated with decreased primary productivity in local waters due to the increased stratification of the California Current off the Oregon coast (Mantua et al. 1997; Hare et al. 1999). These events are likely to affect salmon marine survival rates (Hare et al. 1999).

Changing ocean conditions may affect fish populations through abiotic mechanisms such as modifications to critical habitat. Changes in precipitation regimes associated with the El Niño Southern Oscillation (ENSO), for example, may limit access to spawning grounds, and systems located near the southern end of salmon distribution ranges may reach critical levels as waters warm (Naiman 2002).

Stream Hydrology

Many aspects of climate change are expected to alter the water cycle. The anticipated changes include increased and earlier peak stream flows and reduced summer stream flows (Defenders of Wildlife and ODFW 2008). ODFW (2014) suggests that these changes are likely to compound existing factors that are already limiting the suitability of fish habitat. They recognize the following as factors that are currently affecting critical fish habitat:

- Loss of peripheral stream connections
- Degradation of in-stream structures
- Unfavorable changes to water temperature, sedimentation regimes,

barriers to upstream passage, and availability of gravel.

Increased water temperatures may limit the reproductive function of riverine fish that are already vulnerable. The white sturgeon (*Acipenser transmontanus*), for example, has been shown to have substantial egg mortality in water temperatures above 18 °C (64°F) (Wagoner et al. 1990). Additionally, warmer waters may accelerate habitat loss by effectively eliminating cool backwaters and other areas that provide important refugia and resting habitat for salmon species (ODFW 2014; Defenders of Wildlife and ODFW 2008).

Warmer air temperatures and altered patterns of precipitation are likely to directly influence the frequency, magnitude, and extent of extreme weather including flooding and drought (Reiman and Isaaks 2010; Defenders of Wildlife and ODFW 2008). In instances where these extreme weather events compromise riparian habitat, the effects of climate change on fish may be accelerated by reductions in shading, bank stabilization, food availability, and nutrient and chemical mediation (ODFW 2014).

Literature Cited

- Bakun, A. 1990. Global climate change and intensification of coastal ocean upwelling. *Science*, 247(4939), 198-201.
- Beechie T., H. Imaki, J. Greene, A. Wade, H. Wu, G. Pess, P. Roni, J. Kimball, J. Stanford, P. Kiffney, and N. Mantua. 2012. Restoring Salmon Habitat for a Changing Climate. *River Res. Applic.* Published online in Wiley Online Library (wileyonlinelibrary.com) DOI: 10.1002/rra.2590
- Cornu, C.E., and J. Souder, J. Hamilton, A. Helms, R. Rimler, B. Joyce, F. Reasor, T. Pedersen, E. Wright, R. Namitz, J. Bragg, B. Tanner. 2012. Partnership for Coastal Watersheds State of the South Slough and Coastal Frontal Watersheds. Report prepared for the Partnership for Coastal Watersheds Steering Committee. South Slough National Estuarine Research Reserve and Coos Watershed Association. 225 pp.
- Defenders of Wildlife and Oregon Department of Fish and Wildlife (ODFW). 2008. Preparing Oregon's Fish, Wildlife, and Habitats for Future Climate Change: A Guide for State Adaptation Efforts. Accessed 15 January, 2014. Available online http://www.oregon.gov/ENERGY/GBLWRM/docs/f-w_adaptation_guide.pdf
- Devine, B. M., Munday, P. L., & Jones, G. P. 2012. Homing ability of adult cardinalfish is affected by elevated carbon dioxide. *Oecologia*, 168(1), 269-276.
- Dixon, D. L., Munday, P. L., & Jones, G. P. (2010). Ocean acidification disrupts the innate ability of fish to detect predator olfactory cues. *Ecology Letters*, 13(1), 68-75.
- Environmental Protection Agency (EPA) Office of Water. 2013a. Impacts of Climate Change on the Occurrence of Harmful Algal Blooms. Accessed 4 December, 2013. Available online <http://www2.epa.gov/sites/production/files/documents/climatehabs.pdf>
- Environmental Protection Agency (EPA). 2013b. Climate change indicators in the United States. Retrieved from: <http://www.epa.gov/climatechange/science/indicators/oceans/sea-surface-temp.html>
- Feely, Richard A., Simone R. Alin, Jan Newton, Christopher L. Sabine, Mark Warner, Allan Devol, Christopher Krembs, and Carol Maloy. 2010. The combined effects of ocean acidification, mixing, and

respiration on pH and carbonate saturation in an urbanized estuary. *Estuarine, Coastal and Shelf Science*, 88 (4): 442-449.

Giorgi, A.E., and J. L. Congleton. 1984. Effects of current velocity on development and survival of lingcod, *Ophiodon elongatus*, embryos. *Environmental Biology of Fishes*, 19(1/2):15-27.

Glick, P., J. Clough, and B. Nunley. 2007. Sea-level Rise and Coastal Habitat in the Pacific Northwest: An analysis for Puget Sound, Southwestern Washington, and Northwestern Oregon. National Wildlife Federation. Accessed 15 January 2014. Available online <http://www.nwf.org/pdf/Wildlife/PacificNWSeaLevelRise.pdf>.

Guinotte, J. M., & Fabry, V. J. (2008). Ocean acidification and its potential effects on marine ecosystems. *Annals of the New York Academy of Sciences*, 1134(1), 320-342.

Hamilton, Judy. 2011. SWMP Bio-monitoring: Final Report (2004-2010). South Slough National Estuarine Research Reserve. Charleston, OR.

Hare, S. R., Mantua, N. J., & Francis, R. C. 1999. Inverse production regimes: Alaska and west coast Pacific salmon. *Fisheries*, 24(1), 6-14.

Hayward, T. L. 1997. Pacific Ocean climate change: atmospheric forcing, ocean circulation and ecosystem response. *Trends in ecology & evolution*, 12(4), 150-154.

Ishimatsu, A., Kikkawa, T., Hayashi, M., Lee, K. S., & Kita, J. 2004. Effects of CO₂ on marine fish: larvae and adults. *Journal of oceanography*, 60(4), 731-741.

Kikkawa, T., Ishimatsu, A., & Kita, J. 2003. Acute CO₂ tolerance during the early developmental stages of four marine teleosts. *Environmental Toxicology*, 18(6), 375-382.

Knutson, Thomas R. et al. 2010. Tropical Cyclones and Climate Change. *Nature Geoscience*, 3(3), 1-37.

Landry, C.A., S.L. Steele, S. Manning, A.O. Cheek. 2007. Long Term Hypoxia Suppresses Reproductive Capacity in the Estuarine Fish, *Fundulus granadis*. *Comparative Biochemistry and Physiology*, 148, 317-232.

Mantua, N. J., Hare, S. R., Zhang, Y., Wallace, J. M., & Francis, R. C. 1997. A Pacific interdecadal climate oscillation with impacts on salmon production. *Bulletin of the American Meteorological Society*, 78(6), 1069-1079

McCabe, G. J., M. P. Clark, and M. C. Serreze. 2001. Trends in Northern Hemisphere surface cyclone frequency and intensity, *Journal of Climate*, 14, 2763-2768.

Miller, Jessica A. and Charles A. Simenstad. 1997. A comparative Assessment of a natural and Created Estuarine Slough as Rearing Habitat for Juvenile Chinook and Coho Salmon. *Estuaries*, 20 (4), 792-806.

Monaco, Mark E., and Robert L. Emmett. 1990. Distribution and abundance of fishes and invertebrates in West Coast estuaries. V1 and v2. Rockville, Md: U.S. Dept. of Commerce, National Oceanic and Atmospheric Administration, National Ocean Service.

Munday, P. L., Dixon, D. L., Donelson, J. M., Jones, G. P., Pratchett, M. S., Devitsina, G. V., & Døving, K. B. 2009. Ocean acidification impairs olfactory discrimination and homing ability of a marine fish. *Proceedings of the National Academy of Sciences*, 106(6), 1848-1852.

Mysak LA. 1986. El Niño, interannual variability, and fisheries in the Northeast Pacific Ocean. *Can J Fish Aquat Sci* 43:464-97.

National Research Council. 2012. Sea-Level Rise for the Coasts of California, Oregon, and Washington: Past, Present, and Future. Washington, DC: The National Academies Press. 201 pp.

NOAA Pacific Marine Environmental Laboratory (PMEL) Carbon Program. 2013. What is Ocean Acidification? Retrieved from: <http://www.pmel.noaa.gov/co2/story/What+is+Ocean+Acidification%3F>

Oregon Department of Fish and Wildlife (ODFW). 2006. Strategy Habitat: Riparian Habitats from Oregon Conservation Strategy 2006. Accessed 16 January, 2014. Available online http://www.dfw.state.or.us/conservationstrategy/docs/document_pdf/b-habitat_10.pdf.

Oregon Department of Fish and Wildlife (ODFW). 2014. Coastal Multi-Species Conservation and Management Plan. Unpublished Manuscript.

O'Higgins, Timothy and Steven S. Rumrill. 2007. Tidal and Watershed Forcing of Nutrients and Dissolved Oxygen Stress Within Four Pacific Coast Estuaries: Analysis of Time-Series Data Collected by the National Estuarine Research Reserve System-Wide Monitoring Program (2000-2006) Within Padilla Bay (WA), South Slough (OR), Elkhorn Slough (CA), and the Tijuana River Estuary (CA). Final Report Submitted to the NOAA/Univ. of New Hampshire Cooperative Institute for Coastal and Estuarine Environmental Technology (CICEET). 28 September, 2007.

Pacific Fishery Management Council. 1983. Northern anchovy fishery management plan, fourth draft revision. Pacific Fishery Management Council, Portland, OR.

Perry, Allison L, Paula J. Low, Jim R. Ellis, and John D. Reynolds. 2005. Climate Change and

Distribution Shifts in Marine Fishes. *Science*, 308 (5730), 1912-1915.

Radovich, J. (1961). Relationships of some marine organisms of the northeast Pacific to water temperatures, particularly during 1957 through 1959. *Fish Bull*, 112(62), 18-30.

Rieman, B.E., and D. Isaak. 2010. Climate change, aquatic ecosystems, and fishes in the Rocky Mountain West: implications and alternatives for management. Gen. Tech. Rep. RMRS-GTR-250. Fort Collins, CO: U.S. Department of Agriculture, Forest Service, Rocky Mountain Research Station. 46 p.

Roessig, J. M., Woodley, C. M., Cech Jr, J. J., & Hansen, L. J. 2004. Effects of global climate change on marine and estuarine fishes and fisheries. *Reviews in Fish Biology and Fisheries*, 14(2), 251-275.

Rooper, Christopher N., Donald R. Gunderson, and David A. Armstrong. 2003. Patterns in Use of Estuarine Habitat by Juvenile English Sole (*Pleuronectes vetulus*) in Four Eastern North Pacific Estuaries. *Estuaries*. 26 (4): 1142-1154.

Thomas, P., Rahman, M. S., Khan, I. A., & Kummer, J. A. 2007. Widespread endocrine disruption and reproductive impairment in an estuarine fish population exposed to seasonal hypoxia. *Proceedings of the Royal Society B: Biological Sciences*, 274(1626), 2693-2702.

Wagoner, L.J., K.K. Jones, R.E. Bender, J.A. Butler, D.E. Demory, T.F. Gaumer, J.A. Hurtado, W.G. Mullarky, P.E. Reimers, N.T. Richmond and T.J. Rumreich. 1990. Coos River Basin Fish Management Plan. Oregon Department of Fish and Wildlife, Salem, OR.

Webster, P. J. et al. 2005. Changes in Tropical Cyclone Number, Duration, and Intensity in a Warming Environment. *Science* 309, 1844-1846.

Widdicombe, S., & Spicer, J. I. 2008. Predicting the impact of ocean acidification on benthic biodiversity: what can animal physiology tell us?. *Journal of Experimental Marine Biology and Ecology*, 366(1), 187-197.

Yamanaka, T., D. Raffaelli, and P. White. 2013. Non-linear interactions determine the impact of sea-level rise on estuarine benthic biodiversity and ecosystem processes. *PLoS ONE*, 8(7), 1-8.

Yoklavich, Mary Margaret. 1982. Energy Requirements of the English Sole (*Parophrys vetulus*). Master's thesis. San Francisco State University, San Francisco, CA.